

Trace Inference


Goal: To recover the trace, tangent and curvature fields.

Tangent field: $(x(s), y(s)) \rightarrow (x'(s), y'(s))$

Curvature field: $(x(s), y(s)) \rightarrow (x''(s), y''(s))$

Discretization of Orientation:

θ_λ is the discrete orientation, $\lambda = 1, \dots, m$



$$\theta_\lambda - \pi/2m \leq \theta^* \leq \theta_\lambda + \pi/2m$$

Trace Inference

Let the certainty of tangent with orientation θ_λ at position (x_i, y_i) be:

$$p_i(\lambda) \in [0, 1] \text{ for } i = 1, \dots, n; \lambda = 1, \dots, m$$

The orientation certainty vector is given by:

$$\hat{p}_i = [p_i(1), p_i(2), \dots, p_i(m)]$$

Associate to each orientation vector element $p_i(\lambda)$ a discrete measure of curvature $\kappa_i(\lambda)$.

Trace of Curve?

Singularities?

Hierarchies

Constraints

Quantization

TRACE POINTS

fine



TANGENTS



CURVATURES



CURVATURE CONSISTENCIES



coarse

Two Stages

Stage 1: Measurement

Convolution with linear operators to obtain initial tangent estimates at each position and orientation.

$$G(x, y) = LSF(x) \cdot e^{-y^2/\sigma_y^2}$$
$$LSF(x) = e^{-x^2/\sigma_1^2} - B e^{-x^2/\sigma_2^2} + C e^{-x^2/\sigma_3^2}$$

Stage 2: Interpretation

Threshold to find strongest convolutions?

Discrete Cocircularity

Tangent λ is cocircular to tangent λ' iff

$$\Gamma(\theta, \theta_t) = \Gamma(\theta_t, \theta')$$

for some $\theta, \theta', \theta_t$.

Range of θ is $(\theta_\lambda - \epsilon/2, \theta_\lambda + \epsilon/2)$

Range of θ' is $(\theta_{\lambda'} - \epsilon/2, \theta_{\lambda'} + \epsilon/2)$

Range of θ_t is $(\theta_{ij} - \alpha, \theta_{ij} + \alpha)$

Discrete Cocircularity Condition:

$$|\Gamma(\theta_\lambda, \theta_{ij}) - \Gamma(\theta_{ij}, \theta_{\lambda'})| < \epsilon + 2\alpha$$

Cocircularity support

Measurement stage consists of convolutions against “line detectors” .

With θ_{λ_i} the orientation of the operator at position (x_i, y_i) , the **normalized** convolutions

$$\{p_i(\lambda), i = 1, \dots, n; \lambda = 1, \dots, m\}, 0 \leq p_i(\lambda) \leq 1$$

provide an estimate of the “confidence” in tangent λ at position i .

Cocircularity support for tangent λ at position i :

$$s_i(\lambda) = \sum_{j=1}^n \sum_{\lambda'=1}^m r_{ij}(\lambda, \lambda') p_j(\lambda')$$

where $r_{ij}(\lambda, \lambda') = c_{ij}(\lambda, \lambda')$,
the cocircularity coefficient.

Curvature Classes

Partition the neighborhood support set about tangent A into a discrete set of curvature classes $\mathcal{K}_k(A), k = 1, \dots, K$.

If tangent A is cocircular to B and A is cocircular to C and B, C belong to the **same curvature class** with respect to A , then B is cocircular to C .


Revised cocircularity support function:

$$s_i(\lambda) = \max_{k=1, K} \sum_{j=1}^n \sum_{\lambda'=1}^m r_{ij}^k(\lambda, \lambda') p_j(\lambda')$$

where $r_{ij}(\lambda, \lambda') = c_{ij}(\lambda, \lambda') \cdot K_{ij}^k(\lambda, \lambda')$.


Partitioning Function

$$K_{ij}^k(\lambda, \lambda') = 1 \quad \text{if} \quad \rho_{\min}^k \leq \hat{\rho}_{ij}(\lambda) \leq \rho_{\max}^k \\ = 0 \quad \text{otherwise.}$$



$$\hat{\rho}_{ij}(\lambda) = \frac{d_{ij}}{2 \sin(|\Gamma(\theta_l, \theta_{ij})|)}$$

Curvature Consistency



Modify the consistency coefficients:

$$r_{ij}^{kk'}(\lambda, \lambda') = c_{ij}(\lambda, \lambda') K_{ij}^k(\lambda, \lambda') C_{ij}^{kk'}(\lambda, \lambda')$$

$C_{ij}^{kk'}(\lambda, \lambda') = 1$ if curvature class k of λ is “consistent with curvature class k' of λ' at j ;

$C_{ij}^{kk'}(\lambda, \lambda') = 0$ otherwise.

Average Local Support

$$A(p) = \sum_{i=1}^n s_i(\lambda) p_i(\lambda)$$

The $p_i(\lambda)$'s provide a measure of which tangents are chosen.

The $s_i(\lambda)$'s indicate how mutually consistent they are.

Idea is to iteratively update the $p_i(\lambda)$'s in order to maximize the average local support.
